

Supplementary Information

Synthesis of biologically active 2-thio-5-arylbenzo[4,5]thiazolopyrimido [5,4-d]pyrimidin-4-one derivatives catalyzed by metal proline in water

Medhabati Thiyam, Ranjana Devi Thaodem & Warjeet Singh Laitonjam*

Department of Chemistry, Manipur University, Canchipur, Imphal 795 003, India

E-mail: warjeet@yahoo.com; warjeet@manipuruniv.ac.in

Received 27 January 2020; accepted (revised) 18 August 2021

IR, ^1H NMR and ^{13}C NMR spectra of Metal-proline complexes




Fig. 1 IR spectrum of Cd-Proline Complex




Fig. 2 ¹H NMR spectrum of Cd-Proline Complex




Fig. 3 ¹³C NMR spectrum of Cd-Proline Complex




Fig. 4 IR spectrum of Zn-Proline Complex




Fig. 5 ¹H NMR spectrum of Zn-Proline Complex




Fig.6 ^{13}C NMR spectrum of Zn-Proline Complex




Fig. 7 ^1H NMR spectrum of Hg-Proline Complex




Fig.8 ^{13}C NMR spectrum of Hg-Proline Complex




Fig. 9 IR spectrum of Pb-Proline Complex




Fig. 10 ^1H NMR spectrum of Pb-Proline Complex




Fig. 11 ^{13}C NMR spectrum of Pb-Proline Complex